



Corporate Presentation

Enabling Oral Drug Delivery to
Improve Patient Compliance

May 2020



Forward-Looking Statements

This presentation contains forward-looking statements about Lipocine Inc. (the “Company”). These forward-looking statements are made pursuant to the safe harbor provisions of the Private Securities Litigation Reform Act of 1995. These forward-looking statements relate to the Company’s product candidates, FDA review process related to our resubmitted NDA for TLANDO™, the expected timing of Phase 3 trials for TLANDO XR and LPCN 1107 and Phase 2 studies for LPCN 1144 and LPCN 1148, clinical and regulatory processes and objectives, potential benefits of the Company’s product candidates, intellectual property and related matters, all of which involve known and unknown risks and uncertainties. Actual results may differ materially from the forward-looking statements discussed in this presentation.

Accordingly, the Company cautions investors not to place undue reliance on the forward-looking statements contained in, or made in connection with, this presentation. Several factors may affect the initiation and completion of clinical trials and studies, the potential advantages of the Company’s product candidates and the Company’s capital needs. The forward-looking statements contained in this presentation are qualified by the detailed discussion of risks and uncertainties set forth in the Company’s annual report on Form 10-K and other periodic reports filed by the Company with the Securities and Exchange Commission, all of which can be obtained on the Company’s website at www.lipocine.com or on the SEC website at www.sec.gov. The forward-looking statements contained in this document represent the Company’s estimates and assumptions only as of the date of this presentation and the Company undertakes no duty or obligation to update or revise publicly any forward-looking statements contained in this presentation as a result of new information, future events or changes in the Company’s expectations.

Clinical Stage Biopharmaceutical Company

Innovative Product Candidates for Metabolic and Endocrine Disorders

Proprietary Drug Delivery Platform	PRODUCT (Indication)	PRE-CLINICAL	PHASE 1	PHASE 2	PHASE 3	NDA
	TLANDO™ (Oral Testosterone for Testosterone Replacement Therapy “TRT”)					PDUFA Date August 28, 2020
	TLANDO XR (Long Acting Oral Testosterone for Testosterone Replacement Therapy “TRT”)				Phase 3 Dose Identified	
	LPCN 1144 (Oral Testosterone for Pre-Cirrhotic NASH)			LiFT Phase 2 Clinical Study Ongoing		
	LPCN 1148 (Oral Testosterone for Cirrhosis)			Next Step: POC Phase 2 Clinical Study		
	LPCN 1107 (Oral HPC for Prevention of PTB)				Phase 3 Dose Identified	



TLANDO™

Fixed Dose Oral TRT




Hypogonadism Affects Up to 20M Men^{1, 2}

TLANDO Franchise has the Potential to Drive Market Expansion^{1,2}



Monthly TRT TRx Trend

TRT Market is Growing



2019 Annual TRx of 7.3 million

Issues with Current Non-Oral TRT Options


Potential Barrier To Newly Diagnosed and Existing Patients



- **Black Box Warning**
 - Secondary exposure to testosterone
 - Pulmonary oil micro embolism (POME) and anaphylaxis shock
- **Inconvenient application or painful injection**
- **Poor persistence reflects need for oral**
 - Average days on therapy is 100 days
- **More than 50% of patients need dosage adjustment**
 - Burdensome for patients due to multiple doctor visits


TLANDO Has Potential to Improve Persistence

Titration Requirements of Current TRTs



Number of Current TRT Dose Adjustments by Form*

* Current TRT n=412
Q16. Since you started using your current testosterone medication, how many times was the dose adjusted up or down until you reached your current dose level?



Patient Group	% Persistent Patients		
	Day 30	Day 180	Day 360
New	50.8%	18.0%	9.2%
Experienced	73%	36.7%	21.9%

Cohort Period: February 2016 – January 2017

Analysis Period: 12 Months

Look Back: 6 months for new patients

Source: Adheris Health 2017

TLANDO™ Attributes

Fixed Dose Oral TRT Option

Convenient Oral Route

- Patient and Physician preferred

Easy to Prescribe Fixed Dosing Regimen

- No additional dose adjustment visits
- Less prone to drop out after first Rx
- The “right” dose from the start of therapy with TLANDO™
- Not prone to titration decision errors

Differentiated Hypertension (“HTN”) and Hematocrit Profile

- ~ 1% new anti-HTN starts or increase in anti-HTN dose
- Low incidence of hematocrit increase (erythrocytosis)


Consistent Inter-Day Restoration of T Levels

Demonstrated Paradigm Shifting Liver Benefits



TLANDO Regulatory Update

Near Term PDUFA Date




LPCN 1144

for Non-Cirrhotic NASH



LPCN 1144: Rationale to Target Non-Cirrhotic NASH

Currently No Approved Treatment



Multi-billion \$ Opportunity*

LPCN 1144: Oral Testosterone Therapy

Differentiated NASH Treatment Candidate

Targets Unmet Need

NASH resolution and/or fibrosis improvement

Acceptable tolerability for chronic use

Improvement of sarcopenia

Improvement of sexual dysfunction

Improvement of mental health

Potential Mode of Action

Anti-steatosis

Anti-inflammatory

Anti-oxidative

Cell regenerative

Clinical Experience

Meaningfully reduced liver fat in POC study


Well tolerated in 700+ subjects with up to 52-week exposure

Improved sexual and mood dysfunction

Association Between Testosterone and Liver Disease


Clinical Evidence

NASH



- ~75% of biopsy-confirmed NASH male patients have testosterone < 372 ng/dL¹
- Levels of free T decreased significantly with the increased incidence of fibrosis²

NASH Cirrhosis




“Low T reported in up to 90% of NASH cirrhosis patients³ and is a predictor of mortality.”⁴

LPCN 1144 Proposed Mechanism

Across the Full Spectrum of NASH Pathogenesis

Healthy Liver



Comorbidities

Obesity

Insulin Resistance

Dyslipidemia

Metabolic Syndrome

Hypogonadism (Low T)

LPCN 1144

Anti-steatosis

Steatosis

Anti-inflammatory¹

Inflammation

Anti-oxidant²

Oxidative Stress

Regeneration Booster³

NASH Fibrosis



Removal of Pro-fibrotic inputs⁴



LPCN 1144: Liver Fat Imaging Study (“LFS”)

Study Design and Baseline Liver Fat Subject Distribution



LFS was an open-label, multi-center single-arm 16-week study (N=36) with 225 mg BID LPCN 1144 in hypogonadal males



LF = liver fat
BL = baseline

LPCN 1144: Liver Fat Reduction


Meaningful Relative Liver Fat % Change and Responder Rate



LPCN 1144: Reduction of Liver Injury Markers

In Patients with Elevated ALT at Baseline

Liver enzymes mean change from baseline to end of study (1-Yr), NCT02081300



Alanine amino transferase (ALT), Aspartate amino transferase (AST), Alkaline phosphatase (ALP), Gamma-glutamyl transpeptidase (GGT)

* Sanyal et al, Hepatol, 2015

LPCN 1144: *LiFT* Study* Ongoing

Liver Fat Intervention with Oral Testosterone Study

Phase 2 paired-biopsy clinical study in NASH subjects (NCT04134091)

Study Design


- Three-arm (1:1:1 randomization, two treatments and placebo), multi-center, double-blind
- 225mg twice daily (450mg Daily)
- 20-25 biopsy confirmed NASH male subjects per arm with NAS $\geq$ 4, F1-F3
- Treatment duration of 36 weeks

Primary Endpoint

- Change in hepatic fat fraction via MRI-PDFF measure

Secondary Endpoints

- Change in NASH activity and fibrosis via liver biopsy scoring
- Change in liver enzymes, anthropometric measure, lipids, insulin resistance, inflammatory/fibrosis markers, and labs
- Change in quality-of-life degree (SF-36 and PDQ), weight, BMI, waist circumference, waist to hip ratio, and PAQ activity




LPCN 1148

for Liver Cirrhosis



LPCN 1148: Oral T for Cirrhosis

Background: Prevalence and Common Causes



Common Causes³

Alcoholic liver disease

Nonalcoholic Fatty Liver Disease (NAFLD)

Chronic hepatitis B

Chronic hepatitis C

Cryptogenic

Reduced Testosterone* – Cirrhosis Association

Progressive Drop in T Level with Increasing Disease Severity¹

Increased risk of
major infections,
death and/or
transplantation rates¹

Worsening of
sarcopenia³

Increased risk of for
hepatic
decompensation³

Severity of portal
hypertension and
ascites³

Potential Key Effects of LPCN 1148 in Cirrhotic Patients

Broad Potential Impact

Anabolic	Increase muscle mass and strength Reduce fat mass	Increase bone density Inhibit myostatin	Improve nutritional status*
Androgenic	Induce hematopoiesis Improve sexual dysfunction		
Anti-Inflammatory	Reduce IL-1, IL-6, and TNF- α		
Improve immuno- dysregulation	Lower infection rate		
SHBG lowering	Increase free T		

References: Trivedi and Tapper, Gastroenterol Rep (Oxf), 2018; Berzigotti et al., Hepatology, 2017; Chen and Dunn, Clin Liver Dis (Hoboken), 2018; Sinclair et.al, Liver international, 2016; Neff et al., Digestive Diseases and Sciences, 2004; Puliyl et al., Australian and New Zealand Journal of Medicine, 1977; Brown et al., Cleve Clin Q, 1960; Girolami M, Am Geriatr Soc, 1958; Neff et al., Transplant Proc, 2004; Wells R., The Lancet, 1960; Yurci et al., Clinics and Research in Hepatology and Gastroenterology, 2011; Muting D., Verh Dtsch Ges Inn Med, 1969; Gluud C., Liver, 1984.

*individual's health condition as it is influenced by the intake and utilization of nutrients

Upcoming Milestones

Near Term Value Drivers

	Event	Expected Timing
TLANDO™	PDUFA Date	August 28, 2020
LPCN 1144	<i>LiFT</i> Primary Endpoint Results	4Q 2020

Key Financial Metrics

Stock Price, Market Cap, Cash Balance

Ticker Symbol	LPCN (Nasdaq Capital Market)
Closing Stock Price (5/8/20)	\$0.66/share
Market Capitalization (5/8/20)	\$31.8 million
Cash Balance (3/31/20)	\$20.6 million*
Bank Debt (3/31/20)	\$6.4 million

Lipocine Investment Highlights

Near Term TLANDO PDUFA with a Promising Pipeline

Potential to be a TRT Market Leader

TLANDO™: Differentiated product profile with potential for market expansion

TLANDO XR: Unique long acting oral with potential to maintain leadership

- ~\$2B+ opportunity in an established and growing market with favorable market dynamics

Oral Testosterone Targeted for Pre-Cirrhotic NASH/Cirrhosis

LPCN 1144: A differentiated modality with potential for mono/combo pre-cirrhotic NASH therapy

LPCN 1148: Targeting cirrhosis