

LIPOCINE[®]

ENHANCING HEALTH

Enabling Oral Drug Delivery to Improve Patient Compliance

Corporate Presentation

July 23, 2019



Forward-Looking Statements

This presentation contains forward-looking statements about Lipocine Inc. (the “Company”). These forward-looking statements are made pursuant to the safe harbor provisions of the Private Securities Litigation Reform Act of 1995. These forward-looking statements relate to the Company’s product candidates, FDA review process related to our resubmitted NDA for TLANDO™, the expected timing of Phase 2 studies for LPCN 1144 and LPCN 1148, clinical and regulatory processes and objectives, potential benefits of the Company’s product candidates, intellectual property and related matters, all of which involve known and unknown risks and uncertainties. Actual results may differ materially from the forward-looking statements discussed in this presentation.

Accordingly, the Company cautions investors not to place undue reliance on the forward-looking statements contained in, or made in connection with, this presentation. Several factors may affect the initiation and completion of clinical trials and studies, the potential advantages of the Company’s product candidates and the Company’s capital needs. The forward-looking statements contained in this presentation are qualified by the detailed discussion of risks and uncertainties set forth in the Company’s annual report on Form 10-K and other periodic reports filed by the Company with the Securities and Exchange Commission, all of which can be obtained on the Company’s website at www.lipocine.com or on the SEC website at www.sec.gov. The forward-looking statements contained in this document represent the Company’s estimates and assumptions only as of the date of this presentation and the Company undertakes no duty or obligation to update or revise publicly any forward-looking statements contained in this presentation as a result of new information, future events or changes in the Company’s expectations.

Clinical Stage Biopharmaceutical Company

Metabolic and Endocrine Focus

	PRODUCT (Indication)	PRE-CLINICAL	PHASE 1	PHASE 2	PHASE 3	NDA
Proprietary Drug Delivery Platform	TLANDO (Oral Testosterone for Testosterone Replacement Therapy "TRT")					PDUFA November 9, 2019
	TLANDO XR (LPCN 1111) (Long Acting Oral Testosterone for TRT)				Phase 2 Complete	
	LPCN 1144 (Oral Testosterone for pre-cirrhotic NASH)			LiFT Phase 2 Paired Biopsy Clinical Study in NASH Initiated		
	LPCN 1148 (Oral Testosterone for NASH Cirrhosis)			POC Study Planned		
	LPCN 1107 (Oral HPC for Prevention of PTB)				EOP2 meeting Completed	

TLANDO™ Franchise Status

Differentiated Oral Testosterone Replacement Candidates

TLANDO: PDUFA Date:
November 9, 2019

TLANDO XR
Once-a-Day
Phase 2 Complete

Not subject to any 3rd party data exclusivity




TLANDO™

Fixed Dose Oral Product Targeted for Testosterone Replacement
Therapy


Hypogonadism Affects Up to 20 MM American Men^{1,2}

TLANDO™ Has The Potential To Drive Market Expansion



Monthly TRT TRx Trend

TRT Market is Growing



Source: IMS database
TRx = Total prescriptions

Issues with Current Non-oral TRT Options


Potential Barrier To Newly Diagnosed and Existing Patients



- **Black Box Warning**
 - Secondary exposure to testosterone
 - Pulmonary oil micro embolism (POME) and anaphylaxis shock
- **Inconvenient application or painful injection**
- **Poor persistence reflects need for oral**
 - Average days on therapy is 100 days
- **More than 50% of patients need dosage adjustment**
 - Burdensome for patients due to multiple doctor visits

Discontinuation Rates as High as 50% After Only 30 Days with Topical TRT

TLANDO has Potential to Expand the Market Through Improved Persistence




Patient Group	% Persistent Patients		
	Day 30	Day 180	Day 360
New	50.8%	18.0%	9.2%
Experienced	73.0%	36.7%	21.9%

Cohort Period: February 2016 – January 2017
Analysis Period: 12 Months
Look Back: 6 months for new patients

Source: Adheris Health 2017

Current TRT Dosing Regimes

Over 50% of Patients Require Dose Adjustment with Current TRT Therapies



Number of Current TRT Dose Adjustments by Form*

* Current TRT n=412

Q16. Since you started using your current testosterone medication, how many times was the dose adjusted up or down until you reached your current dose level?

TLANDO Attributes

Patient and Physician Preferred Oral Option

Key Advantages of Oral Route:

- No risk of accidental T transference
- Non-invasive; easy to use
- Less cumbersome/burdensome
- Potential for higher persistence/adherence

PCP preferred fixed dosing regimen

- 450mg (225 mg BID) Daily TU capsule orally administered
 - Unlike most TRT products, fixed “right” dose from the start of therapy with TLANDO for all patients
- Easy to use for patients and physicians to prescribe
- Fewer doctor visits (no dose adjustment visits for patients)
 - More compliant-generalizable to real world male patient behavior
- Not prone to titration decision errors (typical wrong titration decision error range from 10-20%)-risk of patients stuck on wrong dose
- Less prone to drop out after first RX since due to lack of efficacy in patients needing up titration to be efficacious with titrated TRT products
- Fixed/predictable cost for payers with no titration

Differentiated hypertension (“HTN”) profile

- ~ 1% new anti-HTN starts or increase in anti-HTN dose
- 32% of subjects with baseline sBP >140 mm Hg experienced a decrease to <140 mm Hg with mean change of -3 mm Hg

Consistent inter-day restoration of T levels

TLANDO = Opportunity in a Market Ready for Change

Near Term FDA Decision



~\$2B+ Market Opportunity with growing scripts



Significant unmet patient and physician needs



Differentiated fixed dose product profile



Share of voice all time low; detail sensitive; concentrated call points



TLANDO™ PDUFA date is November 9, 2019

TLANDO XR

First Long Acting Oral TRT

TLANDO XR (LPCN 1111): Profile

Once Daily Differentiated Oral TRT

- Once daily oral testosterone
 - New molecule with associated IP
 - Novel prodrug of testosterone for oral delivery through proprietary drug delivery technology
- TLANDO XR proof of concept established
 - Positive Phase 2b study results in hypogonadal men
 - Once daily oral dose provides T levels in the eugonadal range
- Phase 3 daily dose identified based on multiple dose Phase 2 studies in hypogonadal male
- Next steps:
 - Obtain FDA feedback on Phase 3 clinical study design


LPCN 1144

Targeted for Non-Alcoholic Steatohepatitis (“NASH”)

Non-Alcoholic Fatty Liver Disease (“NAFLD”)


No Approved Product for the Treatment of NAFLD/NASH

Fatty liver is a reversible condition wherein large vacuoles of triglyceride (TG) fat accumulate in liver cells via the process of steatosis



Estimated LPCN 1144 Target Market

Reportedly 75% of NASH Male Patients Have Total T less than 372 ng/dL¹



1. Sarkar et al. Gastroenterology 156(6):S-1258 & Poster Sa1623 Digestive Disease Week 2019
2. Estes et al, Hepatol 2018)
3. About 50% of males are accounted in the whole NASH population.

Unmet Need in Pre-Cirrhotic NASH

Currently No Approved Product to Treat Pre-Cirrhotic NASH

Rationale for Using LPCN 1144 Alone or in Combination to Treat Pre-Cirrhotic NASH

Multi-dimensional mechanism of action/efficacy across the full NASH disease spectrum

Tolerable and suitable for chronic use

Majority of biopsy confirmed NASH adult males have low testosterone¹

Sarcopenia in patients with NAFLD is associated with a higher likelihood of having steatohepatitis or advanced liver fibrosis²

Sexual dysfunction is an underappreciated comorbidity in male NASH patients³

1. Sarkar et al. *Gastroenterology* 156(6):S-1258 & Poster Sa1623 Digestive Disease Week 2019

2. Dorota et al., *Sex Med Rev.* 2019, 1-2.

3. Yu et al. *BMC Gastroenterology* (2018) 18:51 <https://doi.org/10.1186/s12876-018-0776-0>

LPCN 1144: Multidimensional Mechanism of Action

Across the Full Spectrum of NASH Pathogenesis

Homeostasis Modifier^{1, 2}

- Alter lipid, cholesterol, and glucose metabolism
- Reduce visceral abdominal fat
- Modify activity of hepatic lipase, and skeletal muscle/adipose lipoprotein lipase

Anti-inflammatory²/ Antioxidant/Immuno- modulator³

- Restore mitochondrial turnover and normalizes oxygen consumption⁴

Regeneration Booster^{5,6}

- Stimulate satellite cells and myocyte precursor resulting in cell differentiation and myocyte proliferation⁷
- Increases circulating endothelial progenitor cells ("EPC")⁸

Anabolic Agent⁹

- Increase muscle mass, bone density in men with liver disease¹⁰

1. Shen and Shi, Int J Endocrinol, 2015

3. Sinclair et al., J Gastroenterol Hepatol, 2015

5. A. Francavilla et al., Digest Dis Sci, 1989

7. Sinha-Hikim et al., J Clin Endocrinol Metab, 2004

9. Gentile MA et al., J Mol Endocrine, 2010

2. Kelly and Jones, J Endocrinol, 2013

4. Linda Vignozzi et al., University of Florence, IT, unpublished, 2018

6. Vic et al., Hepatol 1982



8. Liao CH et al., Andrology, 2013

10. Sinclair et al., J Gastroenterol Hepatol 2016

LPCN 1144: Liver Fat Imaging Study (“LFS”)

Study Design and Baseline Liver Fat Subject Distribution



LFS was an open-label, multi-center single-arm 16-week study (N=36) with LPCN 1144 in hypogonadal males



LF = liver fat

LPCN 1144: Liver Fat Study Results

Meaningful Relative Liver Fat % Change and Responder Rate



* Responder rate for relative change is % of patients with at least 30% for relative change from baseline.


LPCN 1144: Next Step

Advancing Forward

- *LiFT* (Liver Fat intervention with oral Testosterone) Phase 2 paired-biopsy Phase 2 clinical study in NASH subjects – In progress
 - Study Design
 - Three-arm, double-blind placebo controlled
 - Biopsy confirmed NASH male subjects with NAS ≥ 4
 - Paired biopsy at baseline and EOS (36-weeks)
 - First-patient dosing expected in 3Q 2019

LPCN 1144: A Differentiated Oral NASH Therapy Candidate

Prodrug of Endogenous Testosterone



Upcoming Milestones

Near Term Value Drivers

	Event	Expected Timing
TLANDO™	PDUFA Date	November 9, 2019
LPCN 1144	<i>LiFT</i> Phase 2 Paired Biopsy Clinical Study in NASH Subjects - First Patient Dosed	3Q 2019

Key Financial Metrics

Stock Price, Market Cap, Cash Balance

Ticker Symbol	LPCN (Nasdaq Capital Market)
Closing Stock Price (7/19/19)	\$1.83/share
Market Capitalization (7/19/19)	\$45.2 million
Cash Balance (3/31/19)	\$22.5 million*
Bank Debt (3/31/19)	\$9.2 million

* \$5M restricted and becomes unrestricted upon TLANDO approval

Appendix


TLANDO™

Differentiated Oral Product Targeted for Testosterone Replacement
Therapy

Annual TRx ~7M

TLANDO™: NDA Under Review

Near Term FDA Decision



LPCN 1144

Targeted for Non-Alcoholic Steatohepatitis (“NASH”)



A silent killer that affects 30 million Americans¹

Clinical Relationship Between Testosterone and NAFLD

Across the Full Disease Spectrum


Hepatic Steatosis

“Men with low testosterone are at high risk for NAFLD.”¹





NASH

“Levels of free T decreased significantly with the increased incidence of lobular inflammation, hepatocyte ballooning, NAFLD activity score, and fibrosis.”²




Cirrhosis

“Low T levels in cirrhotic men are associated with the combined outcome of death or transplantation.”³



LPCN 1144: Additional Health Benefits

Observed in Hypogonadal Subjects with Elevated ALT*




SF-36, Short Form-36 (0-100); PDQ, Psychosexual Daily Questionnaire (0-7); * ALT > 40 U/L at Baseline in 52 week SOAR Trial (N=33)

LPCN 1144: Extensive Clinical Safety Database

Demonstrated No Unexpected Risks

- 650+ subjects in 14 studies with up to 52 weeks exposure
- No drug related SAEs
- No deaths or MACE events




LPCN 1144: Liver Fat Study Results

Liver Fat Based Subject Distribution at Each Visit


Longer Therapy Improved Proportion of Subjects with Disease Resolution

Baseline Pre treatment




N=34

8 Week Treatment



N=33

16 Week Treatment



N=34

■ LF < 5% (NAFLD Free)

■ 5% ≤ LF < 8%

■ 8% ≤ LF < 10%

■ LF > 10%


LPCN 1148

For Treatment of NASH Cirrhosis

LPCN 1148: Oral T for NASH Cirrhosis

No FDA Approved Product-Transplant Only Cure

Cirrhotic Liver



US Prevalence

Among NASH population (2015)*:

- Fibrosis grade 4 (cirrhosis) case: 1.3M
- Compensated cirrhosis 1.16M
- Decompensated cirrhosis: 134,400

In 2013, cirrhosis cause mortality was ~38,000**
and consistently twice the rate in males as females **

Cirrhotic Patients Characteristics:

- Increased morbidity and mortality
- Symptoms of hypogonadism: altered hair distribution, anemia, sexual dysfunction, testicular atrophy, muscle wasting, fatigue, osteoporosis, gynecomastia
- Late stage symptoms: jaundice, pruritis, hepatic encephalopathy, ascites, anasarca, GI bleeding

*Estes C. et al., Hepatology, 2018; **Yoon and Chen, National Institute on Alcohol Abuse and Alcoholism; surveillance report, 2016

Low Testosterone Increases Adverse Outcome in Male Cirrhotic Patients

T Levels Fall Progressively with Increasing Disease Severity¹

- Low T reported in up to 90% of NASH cirrhosis patients² and is a predictor of mortality³
- Low T associated with:
 - Increased risk of major infections, death and/ or transplantation rates¹
 - Increased risk of for hepatic decompensation⁴
 - Worsening of sarcopenia⁴
 - Higher Child Pugh score grade⁴
 - Severity of portal hypertension and ascites grade⁴
 - Higher MELD score⁵

1. Sinclair et al., Liver Transplantation, 2016

2. Kim et al., Male Hypogonadism, eds: Winters and Huhtaniemi, 2017

3. Sinclair M. et al., J. of Gastro and Hepatology, 2015

4. Paternostro et al, Hepatol Res 2019;

5. Sinclair et.al, Liver international, 2016

LPCN 1148: NASH Cirrhosis

Oral T Therapy

Potentially help patients survive longer while waiting for a liver transplant

- T levels positively correlate with muscle mass in men and modulates bone density, hemoglobin production, insulin resistance, and immunity , commonly impaired in cirrhosis¹
- Testosterone therapy increased muscle mass in men with cirrhosis and low testosterone²
- **Next Steps:**
 - Proof of Concept study in male NASH cirrhosis subjects

1. Sinclair M et al., J. of Gastro and Hepatology, 2016

2. Sinclair M et al., J. of Hepatology, 2016

LPCN 1107

High PTB Medical Costs

≥ \$26 Billion Economic Impact³



- 10% of all US pregnancies¹ (475 -500K) result in PTB (< 37 weeks)-a leading cause of neonatal mortality and morbidity
- First year medical costs for PTB infants are ~ 10x higher than for full term infants²
- 28% of preterm births are to women with histories of early delivery



1. CDC (2016)

2. J. Maternal-Fetal and Neonatal Medicine, Dec. 2006, 19(12), 773-782

3. Institute of Medicine of the National Academies. July 2006

LPCN 1107: Prevention of Preterm Birth (PTB)

United States Market Landscape




\$1B Market Opportunity³

1. AMAG estimates Makena market share based on distributor dispensing data and all other market share based on physician market research data conducted by AMAG.
2. Off guidance represents patients treated outside of guidance of Society for Maternal Fetal Medicine, including patients treated with unapproved therapies and untreated patients.
3. IMS data
4. Annualized September 30, 2018 data

LPCN 1107: First Oral PTB Candidate

Characteristics of the Only Approved Product Options for PTB

Makena 21 gauge needle



IM HPC, Makena®:

- 20-25% patients below reported better efficacy HPC level threshold
- Total of 18-22 injections
 - Injection location: Upper-outer quadrant of the gluteus maximus
 - Weekly visit to/by health care provider
 - ~35% of patients experienced injection site pain during clinical trial
 - ~17% of patients reported site swelling-much greater than placebo during clinical trial

SubQ HPC, Makena®:

- 20-25% patients below reported better efficacy HPC level threshold
- Total of 18-22 injections
 - Approved February 14, 2018
 - Auto injector-ready to use device
 - Injection location: Upper back of the arm
 - Weekly visit to/by health care provider
 - 37.3% of subjects identified injection site pain as a treatment emergent adverse event compared to only 8.2% of subjects in the IM arm

LPCN 1107: First Oral PTB Candidate

Addresses Unmet Need



LPCN 1107- Oral HPC

- Potential for superior efficacy with Phase 3 target dose
- No patient discomfort upon administration
- Steady state achieved in 7 days
- Orphan drug designation
 - Major contribution to patient care
- Next steps:
 - Explore partnering opportunities

LPCN 1107: Economic Impact

Potential Lower PTB Rate – US and Resulting Savings

Assuming 4.3% lower PTB rate relative to Makena[®]



~6000 fewer annual PTBs‡



Estimated annual cost saving in ~\$310M‡‡

‡: Assuming 100% of 140,000 eligible US population treated

‡‡: Assuming ~\$51,600 medical costs/PTB

LPCN 1107: First Oral PTB Candidate

Commercial Outlook/Drivers

First Oral HPC for Prevention of Recurrent PTB

- Preferred route-of-administration is oral

Strong Exclusivity Position

- Orphan Drug Designation
- Technology/IP protection

Potential for Superior Efficacy


- Fewer PTB babies with significant healthcare cost savings

Strong Pharmaco-Economic Justification

- Minimize travel related cost/time and healthcare provider cost/time
- Premium pricing potential to generic IM injections

LPCN 1107: HPC PK-PD Correlation

HPC Concentration and PTB Rate with IM HPC, Makena¹



- Lower % PTB rate can be expected with daily C_{avg} ² HPC levels ≥ 8.2 ng/mL


1. Caritis et al., Am J Obstet Gynecol. 2014 (N=315 subjects)

2. $C_{trough} \cong C_{avg}$ for IM Makena[®]

LPCN 1107: Dose Finding Study Design


PK Study: Oral LPCN 1107 vs IM HPC, Makena

- Open-label, four-period, four-treatment study
- 12 healthy pregnant women- Ages 18-35 years; 16-18 weeks gestation
- All subjects received all four treatments



LPCN 1107: Dose-Finding PK Study Results¹

Oral LPCN 1107 vs IM HPC, Makena



- HPC levels below 8.2 ng/mL:
 - Target LPCN 1107 Phase 3 dose was 0% vs 20% subjects using IM HPC Makena per label
- Average HPC levels at target LPCN 1107 Phase 3 dose
 - ~ 3x greater than the comparator, IM HPC, Makena