



Corporate Presentation

Enabling Oral Drug Delivery to
Improve Patient Compliance

January 2021



Forward-Looking Statements

This presentation contains forward-looking statements about Lipocine Inc. (the “Company”). These forward-looking statements are made pursuant to the safe harbor provisions of the Private Securities Litigation Reform Act of 1995. These forward-looking statements relate to the Company’s products and product candidates, FDA’s approval of TLANDO™, the expected timing of Phase 3 trials for TLANDO XR and LPCN 1107 and Phase 2 studies for LPCN 1144 and LPCN 1148, clinical and regulatory processes and objectives, potential benefits of the Company’s product candidates, intellectual property and related matters, all of which involve known and unknown risks and uncertainties. Actual results may differ materially from the forward-looking statements discussed in this presentation.

Accordingly, the Company cautions investors not to place undue reliance on the forward-looking statements contained in, or made in connection with, this presentation. Several factors may affect the initiation and completion of clinical trials and studies, the potential advantages of the Company’s product candidates and the Company’s capital needs. The forward-looking statements contained in this presentation are qualified by the detailed discussion of risks and uncertainties set forth in the Company’s annual report on Form 10-K and other periodic reports filed by the Company with the Securities and Exchange Commission, all of which can be obtained on the Company’s website at www.lipocine.com or on the SEC website at www.sec.gov. The forward-looking statements contained in this document represent the Company’s estimates and assumptions only as of the date of this presentation and the Company undertakes no duty or obligation to update or revise publicly any forward-looking statements contained in this presentation as a result of new information, future events or changes in the Company’s expectations.

Clinical Stage Biopharmaceutical Company

Innovative Product Candidates for Metabolic and Endocrine Disorders

Proprietary Drug Delivery Platform	PRODUCT (Indication)	PRE-CLINICAL	PHASE 1	PHASE 2	PHASE 3	NDA
	TLANDO™ (Oral Testosterone for Testosterone Replacement Therapy "TRT")					Tentative Approval
	LPCN 1144 (Oral Testosterone for Non-Cirrhotic NASH)			LiFT Phase 2 Clinical Study Ongoing		
	TLANDO XR (Long Acting Oral Testosterone for Testosterone Replacement Therapy "TRT")				Next Step: Food Effect Study	
	LPCN 1148 (Oral Testosterone for Cirrhosis)			Next Step: POC Phase 2 Clinical Study		
	LPCN 1107 (Oral HPC for Prevention of PTB)				Phase 3 Dose Identified	


TLANDO™

The Convenient TRT
without Titration
Requirement



Monthly TRT TRx Trend

TRT Market is Growing



Last 12 months (Nov 2019 – Oct 2020) TRx of 7.7 million

Issues with Current TRT Options


Potential Barriers To Newly Diagnosed and Existing Patients



- **Black box warning**
 - Secondary exposure to testosterone
 - Pulmonary oil micro embolism (POME) and anaphylactic shock
- **Inconvenient application or painful injection**
- **High incidence of erythrocytosis was reported in patients on non-oral**
 - 67% on injectable T, 35% on pellets*
- **Most require dose titrations** - Majority of patients are not on efficacious dose at start of therapy and require up titration
 - Avg titration time to reach efficacious dose takes 3-6 months
 - Finding appropriate dose through titration is burdensome for physicians and patients
 - Requires additional clinic/pharmacy visits and invasive samplings
- **Poor persistence with products requiring titration**
 - 50-80% drop off in 3-6 months
 - Insufficient T levels at the beginning of therapy is one of the top reasons patients stop using TRT

Most TRTs Require Dose Titration in Majority of Patients


Majority of Patients are Not on Efficacious Dose at the Start of Therapy



Number of Current TRT Dose Adjustments by Form*

* Current TRT n=412

Q16. Since you started using your current testosterone medication, how many times was the dose adjusted up or down until you reached your current dose level?




- 74% of patients required dose adjustments
- 97% of dose adjusted patients required up titration

**Jatenzo CLAR-15012 clinical study results

Physician Research Data*

Typical Titration for TRT Takes 3-6 Months to Reach Efficacious Dose for Majority of Patients

Average titration time to steady state dose
N = 402 physicians; 150 primary care, 102 urology, 150 endocrinology



1.18: How long, typically, does it take to titrate a patient to their steady state dose?

TLANDO: De-Risked Regulatory Profile

Eligible to Launch After March 27, 2022

Current Status

- Tentative approvals are clearances for drugs to be marketed, but for the existence of patents or market exclusivities
- Initial launch supplies manufactured
- Business development efforts in progress

Opportunities

- Post COVID pandemic launch
- Adequate time for launch preparation/partnering
- Improves TLANDO XR regulatory prospects


LPCN 1144

for Non-Cirrhotic NASH



LPCN 1144: Rationale to Target Non-Cirrhotic Male NASH Patients


Currently No Approved Treatment



Association Between Testosterone and Liver Disease – Clinical Evidence

Liver Disease in Men Undergoing Androgen Deprivation Therapy (ADT) for Prostate Cancer

Patients Receiving ADT were Significantly More Likely to Have



LPCN 1144: Androgen Receptor Agonist

Differentiated NASH Treatment Candidate

Targets Unmet Need

Acceptable benefit to risk ratio

- NASH resolution and/or fibrosis improvement
- Tolerability for chronic use

Improvement of sarcopenia

Improvement of sexual/mood dysfunction

Clinical Experience

Non-invasive dosing regimen


Meaningfully reduced liver fat in POC study

No notable tolerability issues

Improved sexual/mood dysfunction


LPCN 1144: Androgen Receptor Agonist

Potential Mode of Action



LPCN 1144: Liver Fat Reduction

Meaningful Relative Liver Fat % Change and Responder Rate



LPCN 1144: *LiFT* (Liver Fat Intervention) Study* Ongoing

Phase 2 Paired-Biopsy Clinical Study in NASH Subjects (NCT04134091)

Study Design


- Three-arm (1:1:1 randomization, two treatments and placebo), multi-center, double-blind
- Biopsy confirmed male NASH subjects per arm with NAS ≥ 4 , F1-F3
- Treatment duration of 36 weeks

Primary Endpoint

- Change in hepatic fat fraction via MRI-PDFF

Secondary Endpoints

- Change in NASH activity and fibrosis via liver biopsy scoring
- Change in liver markers, anthropomorphic measures (weight, body mass index (BMI), waist to hip ratio, waist circumference), lipids, insulin resistance, inflammatory/fibrosis markers, and labs
- Change in quality-of-life degree (SF-36 and PDQ) and Physical Activity Questionnaire



TLANDO XR

(Testosterone
Tridecanoate)

Once Daily Oral TRT



TLANDO XR: An Innovative Opportunity

Next Generation TRT Option



TLANDO XR is positioned to be the first Oral QD Product



TLANDO XR is clinically differentiated from TLANDO



Patients and physicians prefer once a day oral testosterone



Strong IP (new molecule & proprietary drug delivery technology)



Positive Phase 2b study results

Phase 2b Study: Result Summary

Met Primary and Secondary Endpoints

- Once daily dose for 14 days in an open label, multiple dose PK study in hypogonadal men (n=36 subjects)
 - ✓ Phase 3 dose identified
 - ✓ No Drug related SAEs
 - ✓ Drug Related AEs are Mild to Moderate
- Next step: Food effect study

LPCN 1148

for Liver Cirrhosis



LPCN 1148: Oral T for Cirrhosis

Background: Prevalence and Common Causes

Transplant Only Cure for Liver Cirrhosis
High Economic Burden of a Liver Transplant¹


Liver Cirrhosis in US

Over 600K patients with liver cirrhosis²


45K deaths in 2017³

62% male vs. 38% female on the liver
transplant (LT) waitlist⁴

Healthy Liver



Liver with Cirrhosis



Common Causes⁵

Alcoholic liver disease

Nonalcoholic Fatty Liver Disease (NAFLD)

Chronic hepatitis B

Chronic hepatitis C

Cryptogenic

Unmet Needs For Patients with Liver Cirrhosis

Areas that LPCN 1148 Addresses



Improvement in quality of life while on waiting list



Stay on transplant list

Sarcopenia is a leading contributory cause of removal from the waitlist


- 40 -70% of cirrhotic patients are sarcopenic



Improvement of post transplant survival/outcomes

Sarcopenia in Liver Cirrhosis

Equivalent to Adding Significant Points to the MELD Score



References: Kim and J.W. Jang, World Journal of Gastroenterology, 2015; Sinclair et al., Journal of Gastroenterology and Hepatology (Australia), 2016; Moctezuma-Velazquez et al., Clinical Nutrition, 2018.; Sinclair et al., World Journal of Gastroenterology, 2017. Montano-Loza et al., Clinical and Translational Gastroenterology, 2015; Lai, J.C., et al., Hepatology, 2017; Englesbe et al., J Am Coll Surg, 2010.

MELD Score: Model For End-Stage Liver Disease Score

LPCN 1107

Prevention of Preterm Birth



Unmet Medical Need

Preterm Birth is a Significant Public Health Issue in the U.S



- One Preterm birth (PTB) every minute¹
- $\geq$ \$26 billion economic impact²
- 10% of all US pregnancies³
- A leading cause of neonatal mortality and morbidity
- Medical costs for PTB infants are ~10x higher than for full term infants⁴

LPCN 1107: Potential To Be The Standard of Care for Prevention of PTB

Oral Hydroxyprogesterone Caproate (HPC)

Convenient dosing

No injection site reactions

First oral enabling technology

Solid IP coverage

Orphan drug designated

Next Step: Pivotal study agreement w/ FDA



Impact of Potential Market Withdrawal of Makena and Its Generics

An Opportunity for LPCN 1107



No approved standard of care for prevention of PTB



Improved market potential



Enriched target population: Not losing high risk patients to active control (injectable IM)



May not require a non-inferiority study; Easier to recruit patients for placebo-controlled trial(s)



Advanced “Phase 3 ready” oral candidate targeting PTB

Upcoming Milestones

Near Term Value Drivers

	Event	Expected Timing
TLANDO™	Full Approval	March 28, 2022
LPCN 1144	<i>LiFT</i> Primary Endpoint Results	January 2021
	<i>LiFT</i> Biopsy Results	Mid 2021

Key Financial Metrics

Stock Price, Market Cap, Cash Balance

Ticker Symbol	LPCN (Nasdaq Capital Market)
Closing Stock Price (12/28/20)	\$1.39/share
Cash Balance (9/30/20)	\$23.8 million
Bank Debt (9/30/20)	\$6.4 million

Appendix

TLANDO™ Attributes*

Convenient Oral Route

- No inadvertent transference or Pulmonary Oil Micro Embolism risks
- Single strength and dose

TRT without titration requirement

- Enables selection of an effective dose at the start of therapy without delay
- No “efficacy gap” upon switching from other TRTs
- No additional pharmacy and clinic copays to reach efficacious dose
- No dose adjustment clinic and pharmacy visits
- No dose adjustment invasive samplings
- No titration decision errors

Bioequivalent exposure in low/med/high fat food

Not Known to Produce Hepatic Adverse Events Associated with 17-methylated Testosterone

*Pending final approval





Physician Research: Physicians View No Titration Product as Positive

- Cited “easy/less titration” as an important advantage of TLANDO™
- Finding the adequate TRT dose through titration is burdensome for physicians and patients

LPCN 1144: Liver Fat Imaging Study (“LFS”)

Study Design and Baseline Liver Fat Subject Distribution


LFS was an open-label, multi-center single-arm 16-week study (N=36) of LPCN 1144 225 mg BID in hypogonadal males



LF = liver fat
BL = baseline

LPCN 1148 in Cirrhotic Patients

Potential Key Effects



References: Trivedi and Tapper, Gastroenterol Rep (Oxf), 2018; Berzigotti et al., Hepatology, 2017; Chen and Dunn, Clin Liver Dis (Hoboken), 2018; Sinclair et.al, Liver international, 2016; Neff et al., Digestive Diseases and Sciences, 2004; Puliyel et al., Australian and New Zealand Journal of Medicine, 1977; Brown et al., Cleve Clin Q, 1960; Girolami M, Am Geriatr Soc, 1958; Neff et al., Transplant Proc, 2004; Wells R., The Lancet, 1960; Yurci et al., Clinics and Research in Hepatology and Gastroenterology, 2011; Muting D., Verh Dtsch Ges Inn Med, 1969; Gluud C., Liver, 1984.

*individual's health condition as it is influenced by the intake and utilization of nutrients

Enabling Oral Drug Delivery to Improve Patient Compliance



LPCN 1107

Prevention of Preterm Births

Preterm Birth (PTB)


Basic Facts

- Risks of adverse outcome for preterm infants rise sharply with decreasing gestation
- Early PTB babies have higher rates of death and disability
 - PTB <35 week has ~6x more relative death risk in first 28 days than full term babies

Gestational Age	Average Length of Stay (Days)	Average Hospital Charges (Dollars)
All Admissions	13.2	\$76,164
<32 weeks	46.2	\$280,811
32-33 weeks	20.3	\$102,182
34-36 weeks	9.8	\$51,083
37-38 weeks	5.9	\$37,137
39-41 weeks	4.9	\$29,771
42+ weeks	6.5	\$47,282

Length of Stay and Hospital Charges Among Special Care Nursery Admissions

Source: National Perinatal Information System/Quality Analytic Services; www.npic.org.
From: March of Dimes Perinatal Data Center, 2011



Economic Impact of PTB on the US Health System

*: Behrman et al. Preterm Birth: Causes, Consequences, and Prevention. Washington, DC: The National Academies Press, 2006, 329-354


Status of MAKENA and its Approved Generic Equivalents

CDER Proposes Withdrawal of Approval for Makena on October 5, 2020

- Confirmatory (PROLONG) study failed to verify clinical benefit and FDA concluded that the available evidence does not show that Makena is effective for its approved use
- CDER has issued a notice of opportunity for a hearing to AMAG Pharmaceuticals and approved generics to Makena for an opportunity to comment
- Makena and its approved generics will remain on the market until the manufacturers decide to remove the drugs or the FDA Commissioner mandates their removal
- FDA intend to hold a meeting with experts in obstetrics, neonatal care, and clinical trial design to discuss how to facilitate development of effective and safe therapies to treat preterm birth
- Eight approved ANDAs referenced Makena as the basis of submission. Upon withdrawal of Makena's NDA, FDA would also withdraw all eight of ANDAs for drug products containing hydroxyprogesterone caproate that reference AMAG's NDA as their reference listed drug

Impact of HPC Plasma Concentration on PTB rate


PK-PD Correlation: HPC Concentration and PTB Rate¹



- Lower % (~4.3%) PTB rate can be expected with daily $C_{avg2} \geq 8.2$ ng/mL compared to overall PTB rate
- 25% of MAKENA (IM HPC) patients with 250 mg/week dosing have levels below the desired “threshold critical for efficacy”

Comparative Steady State PK profiles* to 250 mg/week Injectables HPC

LPCN 1107 Target Regimen Results in Higher Target HPC Levels in Majority of Patients



*PK results obtained from post 35 days for weekly IM Injection & post 8 days of BID dosing for oral from the dose finding study